



Heroin withdrawal symptoms vary in each person.

Many people feel symptoms that are similar to the flu, while others may have more extreme withdrawal symptoms.

Symptoms can start as early as 6 -12 hours from the last time of use.

Overdose symptoms to watch for include:

- Slow and shallow breathing
- Clammy skin
- Blue lips or blue fingernails
- Convulsions or seizures
- Coma

Thinking about change?

Talk with your doctor about treatment options. Together, you can set up a plan for a safe way to make a change.

Improving Wellness in Alabama




Engagement • Collaboration • Education

Visit us at vitalalabama.com

Follow us!



#vitalalabama



Because WE CARE, we ask EVERYONE

HEROIN





vitalalabama.com

Heroin

Heroin is a highly addictive narcotic drug that causes dependence.



Heroin is made from morphine, which is a very powerful painkiller. Heroin acts like a painkiller in the body too, but over time, it breaks down the parts of the brain that make people feel pleasure.

Heroin sold on the street is rarely pure. Instead, it is "cut" with other drugs or substances. It is difficult to know the strength of each dose, so users are at a high risk of overdose or death.

Needles and Heroin:

Sharing needles to inject heroin is very risky. When people share needles, they are at risk for getting:

- HIV
- Hepatitis B
- Hepatitis C

Needle use can also cause complications such as an air bubble in the blood, which can cause a stroke.

Effects of Heroin

